

ON THE CHARACTER SPACE OF BANACH VECTOR-VALUED FUNCTION ALGEBRAS

MORTAZA ABTAHI

ABSTRACT. Given a compact space X and a commutative Banach algebra A , the character spaces of A -valued function algebras on X are investigated. The class of natural A -valued function algebras, those whose characters can be described by means of characters of A and point evaluation homomorphisms, is introduced and studied. For an admissible Banach A -valued function algebra \mathcal{A} on X , conditions under which the character space $\mathfrak{M}(\mathcal{A})$ is homeomorphic to $\mathfrak{M}(\mathfrak{A}) \times \mathfrak{M}(A)$ are presented, where $\mathfrak{A} = C(X) \cap \mathcal{A}$ is the subalgebra of \mathcal{A} consisting of scalar-valued functions. An illustration of the results is given by some examples.

Keywords: Commutative Banach algebras, Banach function algebras, Vector-valued function algebras, Vector-valued characters,

MSC(2010): Primary 46J10; Secondary 46J20.

1. Introduction and Preliminaries

Let A be a commutative unital Banach algebra over the complex field \mathbb{C} . Every nonzero homomorphism $\phi : A \rightarrow \mathbb{C}$ is called a *character* of A . Denoted by $\mathfrak{M}(A)$, the set of all characters of A is nonempty and its elements are automatically continuous [13, Lemma 2.1.5]. Consider the Gelfand transform $\hat{A} = \{\hat{a} : a \in A\}$, where $\hat{a} : \mathfrak{M}(A) \rightarrow \mathbb{C}$ is defined by $\hat{a}(\phi) = \phi(a)$. The *Gelfand topology* of $\mathfrak{M}(A)$ is the weakest topology with respect to which every $\hat{a} \in \hat{A}$ is continuous. Endowed with the Gelfand topology, $\mathfrak{M}(A)$ is compact and Hausdorff. By [13, Theorem 2.1.8], an ideal M in A is maximal if and only if $M = \ker \phi$, for some $\phi \in \mathfrak{M}(A)$. For this reason, sometimes $\mathfrak{M}(A)$ is called the *maximal ideal space* of A . For more on the theory of commutative Banach algebras see, for example, [5, 6, 13, 19].

1.1. Function Algebras. Let X be a compact Hausdorff space and $C(X)$ be the Banach algebra of all continuous functions $f : X \rightarrow \mathbb{C}$ equipped with the uniform norm $\|f\|_X = \sup\{|f(x)| : x \in X\}$. A subalgebra \mathfrak{A} of $C(X)$ is called a *function algebra* on X if \mathfrak{A} separates the points of X and contains the constant functions. If \mathfrak{A} is equipped with some complete algebra norm $\|\cdot\|$, then \mathfrak{A} is called a *Banach function algebra*. If the norm $\|\cdot\|$ of \mathfrak{A} is equivalent to the uniform norm $\|\cdot\|_X$, then \mathfrak{A} is called a *uniform algebra*.

Identifying the character space of a Banach function algebra \mathfrak{A} has been always a problem of interest for mathematicians in this field. For every

$x \in X$, the evaluation homomorphism $\varepsilon_x : f \mapsto f(x)$ is a character of \mathfrak{A} , and the mapping $J : X \rightarrow \mathfrak{M}(\mathfrak{A})$, $x \mapsto \varepsilon_x$, imbeds X homeomorphically as a compact subset of $\mathfrak{M}(\mathfrak{A})$. If J is surjective, one calls \mathfrak{A} *natural* [6, Chapter 4]. In this case, the character space $\mathfrak{M}(\mathfrak{A})$ is identical to X . Note that every semisimple commutative Banach algebra A can be considered, through its Gelfand representation, as a natural Banach function algebra on its character space $\mathfrak{M}(A)$.

A relation between the character space $\mathfrak{M}(\mathfrak{A})$ of a Banach function algebra \mathfrak{A} and the character space $\mathfrak{M}(\bar{\mathfrak{A}})$ of its uniform closure $\bar{\mathfrak{A}}$ was revealed in [10] as follows.

Theorem 1.1 (Honary [10]). *The restriction map $\mathfrak{M}(\bar{\mathfrak{A}}) \rightarrow \mathfrak{M}(\mathfrak{A})$, $\psi \mapsto \psi|_{\mathfrak{A}}$, is a homeomorphism if and only if $\|\hat{f}\| \leq \|f\|_X$, for all $f \in \mathfrak{A}$.*

The above result appears to be very useful in identifying the character spaces in a wide class of Banach function algebras. We establish an analogue of this result for vector-valued function algebras in Section 3.

1.2. Vector-valued Function Algebras. Let A be a commutative unital Banach algebra, and let $C(X, A)$ be the space of all A -valued continuous functions on X . Algebraic operations and the uniform norm $\|\cdot\|_X$ on $C(X, A)$ are defined in the obvious way.

Definition 1.2 (c.f. [2, 15]). A subalgebra \mathcal{A} of $C(X, A)$ is called an *A -valued function algebra* on X if (1) \mathcal{A} contains the constant functions $X \rightarrow A$, $x \mapsto a$, for all $a \in A$, and (2) \mathcal{A} separates the points of X in the sense that, for every pair $x, y \in X$ with $x \neq y$, and for every maximal ideal M of A , there exists some $f \in \mathcal{A}$ such that $f(x) - f(y) \notin M$. If \mathcal{A} is endowed with some algebra norm $\|\cdot\|$ such that the restriction of $\|\cdot\|$ to A is equivalent to the original norm of A and $\|f\|_X \leq \|f\|$, for every $f \in \mathcal{A}$, then \mathcal{A} is called a *normed A -valued function algebra* on X . If the given norm is complete, then \mathcal{A} is called a *Banach A -valued function algebra*. If the given norm is equivalent to the uniform norm $\|\cdot\|_X$, then \mathcal{A} is called an *A -valued uniform algebra*. When no confusion can arise, we use the same notation $\|\cdot\|$ for the norm of \mathcal{A} .

Continuing the work of Yood [18], Hausner [9] proved that τ is a character of $C(X, A)$ if, and only if, there exist a point $x \in X$ and a character $\phi \in \mathfrak{M}(A)$ such that $\tau(f) = \phi(f(x))$, for all $f \in C(X, A)$, whence $\mathfrak{M}(C(X, A))$ is homeomorphic to $X \times \mathfrak{M}(A)$. (In this regard, see [1].) We call a Banach A -valued function algebra *natural* if, like $C(X, A)$, its character space is identical to $X \times \mathfrak{M}(A)$. For instance, in Example 4.1, we will see that the A -valued Lipschitz algebra $\text{Lip}(X, A)$ is natural; see also [7], [11]. Natural A -valued function algebras are studied in Section 2.

1.3. Notations and conventions. Throughout, X is a compact Hausdorff space, and A is a *semisimple* commutative unital Banach algebra. The unit element of A is denoted by $\mathbf{1}$, and the set of invertible elements of A is

denoted by $\text{Inv}(A)$. If $f : X \rightarrow \mathbb{C}$ is a function and $a \in A$, we write fa to denote the A -valued function $X \rightarrow A$, $x \mapsto f(x)a$. If \mathfrak{A} is a function algebra on X , we let $\mathfrak{A}A$ be the linear span of $\{fa : f \in \mathfrak{A}, a \in A\}$, so that any element $f \in \mathfrak{A}A$ is of the form $f = f_1a_1 + \cdots + f_na_n$ with $f_j \in \mathfrak{A}$ and $a_j \in A$. Given an element $a \in A$, we use the same notation a for the constant function $X \rightarrow A$ given by $a(x) = a$, for all $x \in X$, and consider A as a closed subalgebra of $C(X, A)$. Since A has a unit element $\mathbf{1}$, we identify \mathbb{C} with the closed subalgebra $\mathbb{C}\mathbf{1}$ of A . Whence every continuous function $f : X \rightarrow \mathbb{C}$ can be considered as the continuous A -valued function $f\mathbf{1} : x \mapsto f(x)\mathbf{1}$. We drop $\mathbf{1}$ using the same notation f for this A -valued function and adopt the identification $C(X) = C(X)\mathbf{1}$ as a closed subalgebra of $C(X, A)$. Finally, for a family \mathcal{M} of A -valued functions on X , a point $x \in X$, and a character $\phi \in \mathfrak{M}(A)$, we set

$$\mathcal{M}(x) = \{f(x) : f \in \mathcal{M}\}, \quad \phi[\mathcal{M}] = \{\phi \circ f : f \in \mathcal{M}\}.$$

2. NATURAL VECTOR-VALUED FUNCTION ALGEBRAS

Let \mathcal{A} be an A -valued function algebra on X . Assume that M is a maximal ideal of A , $x_0 \in X$, and set

$$(2.1) \quad \mathcal{M} = \{f \in \mathcal{A} : f(x_0) \in M\}.$$

The fact that \mathcal{M} is an ideal of \mathcal{A} is obvious. We prove that \mathcal{M} is maximal. Take a function $g \in \mathcal{A} \setminus \mathcal{M}$ so that $g(x_0) \notin M$. Since M is maximal in A , there exist $a \in M$ and $b \in A$ such that $\mathbf{1} = a + g(x_0)b$. Consider b as a constant function of X into A and let $f = \mathbf{1} - gb$. Then $f(x_0) = a \in M$ so that $f \in \mathcal{M}$ and $\mathbf{1} = f + gb$ which means that the ideal of \mathcal{A} generated by $\mathcal{M} \cup \{g\}$ is equal to \mathcal{A} . Hence \mathcal{M} is maximal in \mathcal{A} .

Definition 2.1. An A -valued function algebra \mathcal{A} on X is called *natural* on X , if every maximal ideal \mathcal{M} of \mathcal{A} is of the form (2.1), for some $x_0 \in X$ and $M \in \mathfrak{M}(A)$.

In case $A = \mathbb{C}$, natural A -valued function algebras coincide with natural (complex) function algebras.

Theorem 2.2. *Let \mathcal{A} be an A -valued function algebra on X . If \mathcal{M} is a maximal ideal in \mathcal{A} and $\mathcal{M}(x_0) \neq A$, for some $x_0 \in X$, then*

- (1) $\mathcal{M}(x_0)$ is a maximal ideal of A ;
- (2) $\mathcal{M}(x) = A$ for $x \neq x_0$;
- (3) $\mathcal{M} = \{f \in \mathcal{A} : f(x_0) \in M\}$, where $M = \mathcal{M}(x_0)$.

Proof. It is easily verified that $\mathcal{M}(x_0)$ is an ideal. We show that $\mathcal{M}(x_0)$ is maximal. Assume that $a \notin \mathcal{M}(x_0)$. Then a , as a constant function on X , does not belong to \mathcal{M} . Hence, the ideal of \mathcal{A} generated by $\mathcal{M} \cup \{a\}$ is equal to \mathcal{A} meaning that $\mathbf{1} = f + ag$, for some $f \in \mathcal{M}$ and $g \in \mathcal{A}$. In particular, $\mathbf{1} = f(x_0) + ag(x_0)$ which implies that the ideal of A generated by $\mathcal{M}(x_0) \cup \{a\}$ is equal to A . Hence, $\mathcal{M}(x_0)$ is maximal.

Now, assume that $x \neq x_0$. Since \mathcal{A} separates the points of X (Definition 1.2), for the maximal ideal $\mathcal{M}(x_0)$ in A , there is a function $f \in \mathcal{A}$ such that $f(x) - f(x_0) \notin \mathcal{M}(x_0)$. Define $g(s) = f(s) - f(x)$ so that $g(x_0) \notin \mathcal{M}(x_0)$. This implies that $g \notin \mathcal{M}$. Since \mathcal{M} is maximal, there are $h \in \mathcal{M}$ and $k \in \mathcal{A}$ such that $h + kg = \mathbf{1}$. Hence, $\mathbf{1} = h(x) \in \mathcal{M}(x)$ and $\mathcal{M}(x) = A$. \square

It is proved in [1] that the algebra $C(X, A)$ satisfies all conditions in Theorem 2.2. Therefore $C(X, A)$ is natural.

Corollary 2.3. *Let \mathcal{A} be an A -valued function algebra on X .*

- (1) *The algebra \mathcal{A} is natural if, and only if, for every proper ideal \mathcal{I} in \mathcal{A} , there exists some $x_0 \in X$ such that $\mathcal{I}(x_0) \neq A$.*
- (2) *If \mathcal{I} is an ideal in \mathcal{A} such that $\mathcal{I}(x_0)$ and $\mathcal{I}(x_1)$, for $x_0 \neq x_1$, are proper ideals in A , then \mathcal{I} cannot be maximal in \mathcal{A} .*

The next discussion requires a concept of zero sets. The zero set of a function $f : X \rightarrow A$ is defined as $Z(f) = \{x : f(x) = \mathbf{0}\}$. This concept of zero set, however, is not useful here in our discussion because, in general, the algebra A may contain nonzero singular elements. Instead, the following slightly modified version of this concept appears to be very useful.

Definition 2.4. For a function $f : X \rightarrow A$, the *singular set* of f is defined to be

$$(2.2) \quad Z_s(f) = \{x \in X : f(x) \notin \text{Inv}(A)\}.$$

The following is an analogy of [6, Proposition 4.1.5 (i)].

Theorem 2.5. *Let \mathcal{A} be a Banach A -valued function algebra on X . Then \mathcal{A} is natural if, and only if, for each finite set $\{f_1, \dots, f_n\}$ of elements in \mathcal{A} with $\bigcap_{j=1}^n Z_s(f_j) = \emptyset$, there exist $g_1, \dots, g_n \in \mathcal{A}$ such that*

$$f_1 g_1 + \dots + f_n g_n = \mathbf{1}.$$

Proof. (\Rightarrow) Suppose that \mathcal{A} is natural and, for a finite set $\{f_1, \dots, f_n\}$ in \mathcal{A} , assume that $Z_s(f_1) \cap \dots \cap Z_s(f_n) = \emptyset$. Let \mathcal{I} be the ideal generated by $\{f_1, \dots, f_n\}$. If $\mathcal{I} \neq \mathcal{A}$, then, since \mathcal{A} is natural, by Corollary 2.3, there exists a point $x_0 \in X$ such that $\mathcal{I}(x_0) \neq A$. In particular, the elements $f_1(x_0), \dots, f_n(x_0)$ are all singular in A , which means that $x_0 \in Z_s(f_1) \cap \dots \cap Z_s(f_n)$, a contradiction. Therefore, $\mathcal{I} = \mathcal{A}$ whence there exist $g_1, \dots, g_n \in \mathcal{A}$ such that $f_1 g_1 + \dots + f_n g_n = \mathbf{1}$.

(\Leftarrow) To show that \mathcal{A} is natural, we take a maximal ideal \mathcal{M} of \mathcal{A} and, using Corollary 2.3, we show that $\mathcal{M}(x_0) \neq A$, for some $x_0 \in X$. Assume, towards a contradiction, that, for every $x \in X$, there exists a function $f_x \in \mathcal{M}$ such that $f_x(x) = \mathbf{1}$. Set $V_x = f_x^{-1}(\text{Inv}(A))$. Then $\{V_x : x \in X\}$ is an open covering of the compact space X . So there exist finitely many points $x_1, \dots, x_n \in X$ such that $X \subset V_{x_1} \cup \dots \cup V_{x_n}$. Then $Z_s(f_{x_1}) \cap \dots \cap Z_s(f_{x_n}) = \emptyset$. By the assumption, there exist functions $g_1, \dots, g_n \in \mathcal{A}$ such that $f_{x_1} g_1 + \dots + f_{x_n} g_n = \mathbf{1}$. Hence, $\mathbf{1} \in \mathcal{M}$, which is a contradiction. \square

Let $f \in \mathcal{A}$ and suppose that $Z_s(f) = \emptyset$ so that $f(X) \subset \text{Inv}(A)$. Since the inverse mapping $a \mapsto a^{-1}$ of $\text{Inv}(A)$ onto itself is continuous, the mapping $x \mapsto f(x)^{-1}$, denoted by $\mathbf{1}/f$, is a continuous A -valued function on X . Hence f is invertible in $C(X, A)$. However, f may not be invertible in \mathcal{A} . Let us call \mathcal{A} a *full subalgebra* of $C(X, A)$ if every $f \in \mathcal{A}$ that is invertible in $C(X, A)$ is invertible in \mathcal{A} . The following is an analogy of [3, Theorem 2.1].

Theorem 2.6. *Let \mathcal{A} be a Banach A -valued function algebra on X such that $\bar{\mathcal{A}}$, the uniform closure of \mathcal{A} , is natural. If $\mathbf{1}/f \in \mathcal{A}$ whenever $f \in \mathcal{A}$ and $Z_s(f) = \emptyset$, then \mathcal{A} is natural.*

Proof. We apply Theorem 2.5 to prove that \mathcal{A} is natural. Let f_1, \dots, f_n be elements in \mathcal{A} such that $Z_s(f_1) \cap \dots \cap Z_s(f_n) = \emptyset$. We prove the existence of a finite set $\{g_1, \dots, g_n\}$ of elements in \mathcal{A} such that $f_1g_1 + \dots + f_ng_n = \mathbf{1}$. Regarding f_1, \dots, f_n as elements of $\bar{\mathcal{A}}$, since $\bar{\mathcal{A}}$ is natural, again by Theorem 2.5, there exist h_1, \dots, h_n in $\bar{\mathcal{A}}$ such that $f_1h_1 + \dots + f_nh_n = \mathbf{1}$. For each h_j , there is some $g_j \in \mathcal{A}$ such that $\|h_j - g_j\|_X < \sum_{j=1}^n \|f_j\|_X$. Thus

$$(2.3) \quad \left\| \mathbf{1} - \sum_{j=1}^n f_j g_j \right\|_X = \left\| \sum_{j=1}^n f_j h_j - \sum_{j=1}^n f_j g_j \right\|_X \leq \sum_{j=1}^n \|f_j\|_X \|h_j - g_j\|_X < 1.$$

Hence, for every $x \in X$, $f(x) = \sum f_j(x)g_j(x)$ is an invertible element of A , so that for the function $f = \sum f_j g_j$, which belongs to \mathcal{A} , we have $Z_s(f) = \emptyset$. By the assumption, there is a function g in \mathcal{A} such that $\mathbf{1} = fg = \sum f_j(g_j g)$. Now, Theorem 2.5 shows that \mathcal{A} is natural. \square

An application of the above theorem is given in Example 4.1.

Let \mathcal{A} be a Banach A -valued function algebra. For every point $x \in X$ and character $\phi \in \mathfrak{M}(A)$ define

$$\varepsilon_x \diamond \phi : \mathcal{A} \rightarrow \mathbb{C}, \quad \varepsilon_x \diamond \phi(f) = \varepsilon_x(\phi \circ f) = \phi(f(x)).$$

Then $\varepsilon_x \diamond \phi$ is a character of \mathcal{A} with $\ker(\varepsilon_x \diamond \phi) = \{f \in \mathcal{A} : f(x) \in \ker \phi\}$, which of course is of the form (2.1). Define

$$(2.4) \quad \mathcal{J} : X \times \mathfrak{M}(A) \rightarrow \mathfrak{M}(\mathcal{A}), \quad (x, \phi) \rightarrow \varepsilon_x \diamond \phi.$$

Theorem 2.7. *The mapping \mathcal{J} is a homeomorphism of $X \times \mathfrak{M}(A)$ onto a compact subset of $\mathfrak{M}(\mathcal{A})$. If \mathcal{A} is natural, then $\mathfrak{M}(\mathcal{A})$ is homeomorphic to $X \times \mathfrak{M}(A)$.*

Proof. Take $x \in X$, $\phi \in \mathfrak{M}(A)$, and set $\tau_0 = \varepsilon_x \diamond \phi$. Let W be a neighbourhood of τ_0 in $\mathfrak{M}(\mathcal{A})$ of the form

$$W = \{\tau \in \mathfrak{M}(\mathcal{A}) : |\tau(f_i) - \tau_0(f_i)| < \varepsilon, 1 \leq i \leq n\},$$

where $f_1, \dots, f_n \in \mathcal{A}$. Take

$$U = \{y \in X : \|f_i(y) - f_i(x)\| < \varepsilon/2, 1 \leq i \leq n\},$$

$$V = \{\psi \in \mathfrak{M}(A) : |\psi(f_i(x)) - \phi(f_i(x))| < \varepsilon/2, 1 \leq i \leq n\}.$$

Then U is a neighbourhood of x in X and V is a neighbourhood of ϕ in $\mathfrak{M}(A)$, so that $U \times V$ is a neighbourhood of (x, ϕ) in $X \times \mathfrak{M}(A)$. If $(y, \psi) \in U \times V$ then, for every i ($1 \leq i \leq n$),

$$\begin{aligned} |\psi(f_i(y)) - \phi(f_i(x))| &\leq |\psi(f_i(y)) - \psi(f_i(x))| + |\psi(f_i(x)) - \phi(f_i(x))| \\ &< \|\psi\| \|f_i(y) - f_i(x)\| + \varepsilon/2 \leq \varepsilon/2 + \varepsilon/2 = \varepsilon. \end{aligned}$$

This shows that $\varepsilon_y \diamond \psi \in W$ and thus \mathcal{J} is continuous. Finally, if \mathcal{A} is natural then every maximal ideal of \mathcal{A} is of the form (2.1) which means that every character $\tau \in \mathfrak{M}(\mathcal{A})$ is of the form $\tau = \varepsilon_x \diamond \phi$, for some $x \in X$ and $\phi \in \mathfrak{M}(A)$. Hence, \mathcal{J} is a surjection and thus a homeomorphism. \square

3. CHARACTERS ON VECTOR-VALUED FUNCTION ALGEBRAS

We turn to a more general case where a vector-valued function algebra may not be natural. Let \mathcal{A} be a Banach A -valued function algebra. We show that, under certain conditions, the character space $\mathfrak{M}(\mathcal{A})$ is identical to $\mathfrak{M}(\mathfrak{A}) \times \mathfrak{M}(A)$, where $\mathfrak{A} = C(X) \cap \mathcal{A}$ is the subalgebra of \mathcal{A} consisting of scalar-valued functions. To this end, we should restrict ourself to the class of admissible algebras. If $f \in \mathcal{A}$ and $\phi \in \mathfrak{M}(A)$, it is clear that $\phi \circ f \in C(X)$; it is not, however, clear whether the A -valued function $(\phi \circ f)\mathbf{1}$ belongs to \mathcal{A} . In fact, [2, Example 2.4] shows that it may very well happen that $(\phi \circ f)\mathbf{1} \notin \mathcal{A}$.

Definition 3.1 ([2]). The A -valued function algebra \mathcal{A} is called *admissible* if

$$(3.1) \quad \{(\phi \circ f)\mathbf{1} : f \in \mathcal{A}, \phi \in \mathfrak{M}(A)\} \subset \mathcal{A}.$$

Note that \mathcal{A} is admissible if, and only if, $\phi[\mathcal{A}]\mathbf{1} \subset \mathcal{A}$, for all $\phi \in \mathfrak{M}(A)$.

Admissible vector-valued function algebras exist around in abundant. Some typical examples are $C(X, A)$, $\text{Lip}(X, A)$, $P(K, A)$, $R(K, A)$, etc. Tensor products of the form $\mathfrak{A} \otimes A$, where \mathfrak{A} is a (Banach) function algebra on X , can be seen as admissible A -valued function algebras. (More details are given in Example 4.4.)

During this section, we assume that \mathcal{A} is admissible and set $\mathfrak{A} = \mathcal{A} \cap C(X)$. Then \mathfrak{A} is the subalgebra of \mathcal{A} consisting of all complex functions in \mathcal{A} , it forms a complex function algebra by itself, and $\mathfrak{A} = \phi[\mathcal{A}]$, for all $\phi \in \mathfrak{M}(A)$. Our aim is to give a description of maximal ideals in \mathcal{A} . To begin, take a character $\phi \in \mathfrak{M}(A)$ and a maximal ideal \mathbf{M} of \mathfrak{A} , and set

$$(3.2) \quad \mathcal{M} = \{f \in \mathcal{A} : \phi \circ f \in \mathbf{M}\}.$$

Then \mathcal{M} is a maximal ideal of \mathcal{A} . One way to see this (though it can be seen directly) is as follows. Take $\psi \in \mathfrak{M}(\mathfrak{A})$ with $\mathbf{M} = \ker \psi$ and define

$$\psi \diamond \phi : \mathcal{A} \rightarrow \mathbb{C}, \quad \psi \diamond \phi(f) = \psi(\phi \circ f).$$

Note that $\psi(\phi \circ f)$ is meaningful since $\phi \circ f \in \mathfrak{A}$. The functional $\psi \diamond \phi$ is a character of \mathcal{A} with $\ker(\psi \diamond \phi) = \mathcal{M}$. Hence \mathcal{M} is a maximal ideal of \mathcal{A} .

The main question is whether any maximal ideal \mathcal{M} of \mathcal{A} is of the form (3.2).

Lemma 3.2. *A maximal ideal \mathcal{M} of \mathcal{A} is of the form (3.2) if and only if $\phi[\mathcal{M}] \neq \mathfrak{A}$ for some $\phi \in \mathfrak{M}(A)$.*

Proof. If \mathcal{M} is of the form (3.2) then clearly $\phi[\mathcal{M}] \neq \mathfrak{A}$. Conversely, assume that $\phi[\mathcal{M}] \neq \mathfrak{A}$ for some $\phi \in \mathfrak{M}(A)$. Then $\phi[\mathcal{M}]$ is an ideal of \mathfrak{A} . We show that it is maximal. If $g \notin \phi[\mathcal{M}]$, then $g = g\mathbf{1}$ (as an A -valued function on X) does not belong to \mathcal{M} . Since \mathcal{M} is maximal in \mathcal{A} , the ideal generated by $\mathcal{M} \cup \{g\}$ is equal to \mathcal{A} . This implies that $\mathbf{1} = f + gh$, for some $f \in \mathcal{M}$ and $h \in \mathcal{A}$. Since $\phi \circ g = g$, we get $\mathbf{1} = \phi \circ f + g(\phi \circ h)$. This means that the ideal of \mathfrak{A} generated by $\phi[\mathcal{M}] \cup \{g\}$ is equal to \mathfrak{A} . Thus, $\phi[\mathcal{M}]$ is maximal. Set $\mathfrak{M} = \phi[\mathcal{M}]$ and $\mathcal{M}_1 = \{f \in \mathcal{A} : \phi \circ f \in \mathfrak{M}\}$. Then $\mathcal{M} \subset \mathcal{M}_1$ and both \mathcal{M} and \mathcal{M}_1 are maximal ideals. Hence, $\mathcal{M} = \mathcal{M}_1$. \square

If $\mathcal{M} = \ker \tau$, for some $\tau \in \mathfrak{M}(\mathcal{A})$, then \mathcal{M} is of the form (3.2) if and only if $\tau = \psi \diamond \phi$, for some $\psi \in \mathfrak{M}(\mathfrak{A})$ and $\phi \in \mathfrak{M}(A)$. Let us extend the mapping \mathcal{J} in (2.4) to $\mathfrak{M}(\mathfrak{A}) \times \mathfrak{M}(A)$ as follows.

$$(3.3) \quad \mathcal{J} : \mathfrak{M}(\mathfrak{A}) \times \mathfrak{M}(A) \rightarrow \mathfrak{M}(\mathcal{A}), \quad \mathcal{J}(\psi, \phi) = \psi \diamond \phi.$$

The mapping is injective for if $\psi \diamond \phi = \psi' \diamond \phi'$ then

$$\begin{aligned} \phi(a) &= \psi(\phi(a)) = \psi'(\phi'(a)) = \phi'(a) \quad (a \in A), \\ \psi(f) &= \psi(\phi(f)) = \psi'(\phi'(f)) = \psi'(f) \quad (f \in \mathfrak{A}), \end{aligned}$$

which implies that $\phi = \phi'$ and $\psi = \psi'$. The main question is whether \mathcal{J} is surjective. If $\tau \in \mathfrak{M}(\mathcal{A})$ then $\phi = \tau|_A \in \mathfrak{M}(A)$ and $\psi = \tau|_{\mathfrak{A}} \in \mathfrak{M}(\mathfrak{A})$. The question is whether the equality $\tau = \psi \diamond \phi$ holds true; of course, it does hold if $\phi[\mathcal{M}] \neq \mathfrak{A}$.

Theorem 3.3. *If the mapping \mathcal{J} in (3.3) is a surjection, then it is a homeomorphism and, therefore, the character space $\mathfrak{M}(\mathcal{A})$ is identical to $\mathfrak{M}(\mathfrak{A}) \times \mathfrak{M}(A)$.*

Proof. Suppose that \mathcal{J} is a surjection (and thus a bijection). Since both the domain and the range are compact Hausdorff spaces, it suffices to prove that \mathcal{J} is open. Take $\psi_0 \in \mathfrak{M}(\mathfrak{A})$, $\phi_0 \in \mathfrak{M}(A)$ and set $\tau_0 = \mathcal{J}(\psi_0, \phi_0) = \psi_0 \diamond \phi_0$. Let U and V be neighborhoods of ψ_0 and ϕ_0 of the following form

$$\begin{aligned} U &= \{\psi \in \mathfrak{M}(\mathfrak{A}) : |\psi(f) - \psi_0(f)| < \varepsilon_1 \quad (f \in F_1)\}, \\ V &= \{\phi \in \mathfrak{M}(A) : |\phi(a) - \phi_0(a)| < \varepsilon_2 \quad (a \in F_2)\}, \end{aligned}$$

where F_1 and F_2 are finite sets in \mathfrak{A} and A , respectively. Take $F = F_1 \cup F_2$ as a finite set in \mathcal{A} , $\varepsilon = \min\{\varepsilon_1, \varepsilon_2\}$ and set

$$W = \{\tau \in \mathfrak{M}(\mathcal{A}) : |\tau(f) - \tau_0(f)| < \varepsilon \quad (f \in F)\}.$$

Then W is a neighborhood of τ_0 in $\mathfrak{M}(\mathcal{A})$ and $\mathcal{J}(U \times V) \subset W$. Hence \mathcal{J} is open. \square

The rest of this section is devoted to investigating conditions under which \mathcal{J} is surjective.

Theorem 3.4 (\mathcal{P}). *For a character $\tau \in \mathfrak{M}(\mathcal{A})$ with $\mathcal{M} = \ker \tau$ and $\phi = \tau|_A$, the following are equivalent.*

- (i) $\phi[\mathcal{M}] \neq \mathfrak{A}$.
- (ii) \mathcal{M} is of the form (3.2) with $M = \phi[\mathcal{M}]$.
- (iii) For every $f \in \mathcal{A}$, if $\phi \circ f = \mathbf{0}$ then $f \in \mathcal{M}$.
- (iv) For every $f \in \mathcal{A}$, $\tau(\phi \circ f) = \tau(f)$.
- (v) For every $f \in \mathcal{A}$, if $f(X) \subset \mathcal{M}$ then $f \in \mathcal{M}$.
- (vi) $\tau = \psi \diamond \phi$, for some $\psi \in \mathfrak{M}(\mathfrak{A})$.

Proof. The equivalence (i) \Leftrightarrow (ii) is just Lemma 3.2. The implication (ii) \Rightarrow (iii) is clear. To see the implication (iii) \Rightarrow (iv), let $g = f - (\phi \circ f)\mathbf{1}$. Then $\phi \circ g = \mathbf{0}$ and thus $g \in \mathcal{M}$ and $\tau(g) = 0$. Hence, $\tau(\phi \circ f) = \tau(f)$. The implication (iv) \Rightarrow (iii) is clear.

To prove (iii) \Leftrightarrow (v), we note that $f(X) \subset \mathcal{M}$ if and only if $\phi \circ f = \mathbf{0}$. In fact, $f(X) \subset \mathcal{M}$ means that, for every $x \in X$, the element $f(x)$, as a constant function of X into A , belongs to \mathcal{M} . This, in turn, means that $\tau(f(x)) = \phi(f(x)) = 0$, for all $x \in X$, which means that $\phi \circ f = \mathbf{0}$.

To prove (iii) \Rightarrow (vi), first note that \mathcal{A} being admissible implies that

$$\mathfrak{A} = \phi[\mathcal{A}] = \{\phi \circ f : f \in \mathcal{A}\}.$$

Define $\psi : \mathfrak{A} \rightarrow \mathbb{C}$ by $\psi(\phi \circ f) = \tau(f)$. This is well-defined for if $\phi \circ f = \phi \circ g$ then, by the assumption, $f - g \in \mathcal{M}$ which in turn implies that $\tau(f) = \tau(g)$. Obviously, $\psi \in \mathfrak{M}(\mathfrak{A})$ and $\tau = \psi \diamond \phi$.

Finally, we prove that (vi) \Rightarrow (i). Towards a contradiction, assume that $\phi[\mathcal{M}] = \mathfrak{A}$. Then $\phi \circ f = \mathbf{1}$, for some $f \in \mathcal{M}$. Hence $1 = \psi(\mathbf{1}) = \psi(\phi \circ f) = \tau(f) = 0$ which is absurd. \square

Convention. We say that ' \mathcal{A} has property \mathcal{P} ' if every $\mathcal{M} \in \mathfrak{M}(\mathcal{A})$ satisfies one (and hence all) of conditions in Theorem 3.4. Hence \mathcal{A} has \mathcal{P} if and only if the mapping \mathcal{J} in (3.3) is surjective.

Let $\bar{\mathcal{A}}$ denote the uniform closure of \mathcal{A} in $C(X, A)$. The restriction map

$$(3.4) \quad \mathfrak{M}(\bar{\mathcal{A}}) \rightarrow \mathfrak{M}(\mathcal{A}), \quad \bar{\tau} \mapsto \bar{\tau}|_{\mathcal{A}},$$

is one-to-one and continuous with respect to the Gelfand topology [10]. We write $\mathfrak{M}(\bar{\mathcal{A}}) = \mathfrak{M}(\mathcal{A})$ if it is a homeomorphism.

Proposition 3.5. *If \mathcal{A} has \mathcal{P} then $\bar{\mathcal{A}}$ has \mathcal{P} . If $\bar{\mathcal{A}}$ has \mathcal{P} and $\|f\| \leq \|f\|_X$, for all $f \in \mathcal{A}$, then \mathcal{A} has \mathcal{P} .*

Proof. Suppose that \mathcal{A} has \mathcal{P} . Take $\bar{\tau} \in \mathfrak{M}(\bar{\mathcal{A}})$, set $\tau = \bar{\tau}|_{\mathcal{A}}$ and $\phi = \bar{\tau}|_A = \tau|_A$. Since \mathcal{A} has \mathcal{P} , by Theorem 3.4 (iv), $\tau(\phi \circ f) = \tau(f)$, for all $f \in \mathcal{A}$. Given $f \in \bar{\mathcal{A}}$, there is a sequence $\{f_n\}$ in \mathcal{A} such that $\|f_n - f\|_X \rightarrow 0$. Hence, $\|\phi \circ f_n - \phi \circ f\|_X \rightarrow 0$, and thus

$$\bar{\tau}(\phi \circ f) = \lim_{n \rightarrow \infty} \bar{\tau}(\phi \circ f_n) = \lim_{n \rightarrow \infty} \tau(\phi \circ f_n) = \lim_{n \rightarrow \infty} \tau(f_n) = \lim_{n \rightarrow \infty} \bar{\tau}(f_n) = \bar{\tau}(f).$$

Again, by Theorem 3.4 (iv), we see that $\tilde{\mathcal{A}}$ has \mathcal{P} .

Now, assume that $\tilde{\mathcal{A}}$ has \mathcal{P} , and $\|\hat{f}\| \leq \|f\|_X$, for all $f \in \mathcal{A}$. Take $\tau \in \mathfrak{M}(\mathcal{A})$ and $\phi = \tau|_A$. Extend τ to a character $\bar{\tau} : \tilde{\mathcal{A}} \rightarrow \mathbb{C}$ (this is possible since $\|\hat{f}\| \leq \|f\|_X$, for all $f \in \mathcal{A}$). Note that still we have $\phi = \bar{\tau}|_A$. Since $\tilde{\mathcal{A}}$ satisfies \mathcal{P} , we have $\bar{\tau}(\phi \circ f) = \bar{\tau}(f)$, for all $f \in \tilde{\mathcal{A}}$. This implies that $\tau(\phi \circ f) = \tau(f)$, for all $f \in \mathcal{A}$, and thus \mathcal{A} has \mathcal{P} . \square

The following is a vector-valued version of Theorem 1.1.

Theorem 3.6. *For an admissible Banach A -valued function algebra \mathcal{A} with $\mathfrak{A} = C(X) \cap \mathcal{A}$, let $\tilde{\mathcal{A}}$ and $\tilde{\mathfrak{A}}$ be the uniform closures of \mathcal{A} and \mathfrak{A} , respectively. Consider the following statements:*

- (i) $\mathfrak{M}(\tilde{\mathcal{A}}) = \mathfrak{M}(\mathcal{A})$.
- (ii) $\|\hat{f}\| \leq \|f\|_X$, for all $f \in \mathcal{A}$.
- (iii) $\|\hat{f}\| \leq \|f\|_X$, for all $f \in \mathfrak{A}$.
- (iv) $\mathfrak{M}(\tilde{\mathfrak{A}}) = \mathfrak{M}(\mathfrak{A})$.

Then (i) \Leftrightarrow (ii) \Rightarrow (iii) \Leftrightarrow (iv). If \mathcal{A} satisfies \mathcal{P} , then (iii) \Rightarrow (ii).

Proof. The equivalences (i) \Leftrightarrow (ii) and (iii) \Leftrightarrow (iv) follow from the main theorem in [10]. The implication (ii) \Rightarrow (iii) is obvious, because $\mathfrak{A} \subset \mathcal{A}$.

Assume that \mathcal{A} satisfies \mathcal{P} , and $\|\hat{f}\| \leq \|f\|_X$, for all $f \in \mathfrak{A}$. Fix a function $f \in \mathcal{A}$ and take an arbitrary character $\tau \in \mathfrak{M}(\mathcal{A})$. Since \mathcal{A} has \mathcal{P} , we have $\tau = \psi \circ \phi$, where $\psi = \tau|_{\mathfrak{A}}$ and $\phi = \tau|_A$. Since $\phi \circ f \in \mathfrak{A}$, we have

$$|\tau(f)| = |\psi(\phi \circ f)| \leq \|\widehat{\phi \circ f}\| \leq \|\phi \circ f\|_X \leq \|f\|_X.$$

Hence $\|\hat{f}\| \leq \|f\|_X$, for all $f \in \mathcal{A}$. \square

4. EXAMPLES

To illustrate the results, we devote this section to some examples.

Example 4.1. Let (X, ρ) be a compact metric space. A function $f : X \rightarrow A$ is called an A -valued Lipschitz function if

$$(4.1) \quad L(f) = \sup \left\{ \frac{\|f(x) - f(y)\|}{\rho(x, y)} : x, y \in X, x \neq y \right\} < \infty.$$

The space of A -valued Lipschitz functions on X is denoted by $\text{Lip}(X, A)$. For any $f \in \text{Lip}(X, A)$, the Lipschitz norm of f is defined by $\|f\|_L = \|f\|_X + L(f)$. This makes $\text{Lip}(X, A)$ an admissible Banach A -valued function algebra on X with $\text{Lip}(X) = \text{Lip}(X, A) \cap C(X)$, where $\text{Lip}(X) = \text{Lip}(X, \mathbb{C})$ is the classical complex Lipschitz algebra on X .

The algebra $\text{Lip}(X)$ satisfies all conditions in the Stone-Weierstrass Theorem and thus it is dense in $C(X)$. On the other hand, by [9, Lemma 1], $C(X)A$ is dense in $C(X, A)$ and thus $\text{Lip}(X)A$ is dense in $C(X, A)$. Since $\text{Lip}(X, A)$ contains $\text{Lip}(X)A$, we see that $\text{Lip}(X, A)$ is dense in $C(X, A)$.

It is easy to verify that if $f \in \text{Lip}(X, A)$ and $Z_s(f) = \emptyset$, then $1/f \in \text{Lip}(X, A)$. Since $C(X, A)$ is natural, Theorem 2.6 now implies that $\text{Lip}(X, A)$

is natural. By Theorem 2.7, $\mathfrak{M}(\text{Lip}(X, A))$ is homeomorphic to $X \times \mathfrak{M}(A)$. See also [7] and [11].

Example 4.2. Assume that $A = \mathbb{C}^n$, for some positive integer n . Then, for every admissible Banach A -valued function algebra \mathcal{A} on X , the mapping \mathcal{J} in (3.3) is surjective and thus $\mathfrak{M}(\mathcal{A})$ is identical to $\mathfrak{M}(\mathfrak{A}) \times \mathfrak{M}(\mathbb{C}^n)$.

To see this, we show that \mathcal{A} satisfies condition (i) of Theorem 3.4. Note that $\mathfrak{M}(\mathbb{C}^n) = \{\pi_1, \dots, \pi_n\}$, where $\pi_i : \mathbb{C}^n \rightarrow \mathbb{C}$ is the projection on i -th component. Assume \mathcal{M} is an ideal in \mathcal{A} and $\mathbf{1} \in \pi_i[\mathcal{M}]$, for all $i = 1, \dots, n$. Hence, for every i , there is some $f_i \in \mathcal{M}$ such that $\pi_i \circ f_i = \mathbf{1}$. Let $\{e_1, \dots, e_n\}$ be the standard basis of \mathbb{C}^n . Then e_i , as a constant function of X into A , belongs to \mathcal{A} . Since \mathcal{M} is an ideal, we have $\mathbf{1} = e_1 f_1 + \dots + e_n f_n \in \mathcal{M}$. Hence, $\mathcal{M} = \mathcal{A}$ and \mathcal{M} cannot be maximal.

If $\mathcal{X} = \{1, \dots, n\}$, then $\mathbb{C}^n = C(\mathcal{X})$. The above example states that, given any admissible Banach $C(\mathcal{X})$ -valued function algebra, we have $\mathfrak{M}(\mathcal{A}) = \mathfrak{M}(\mathfrak{A}) \times \mathfrak{M}(C(\mathcal{X}))$. If \mathcal{X} is an arbitrary compact Hausdorff space, it is unknown whether the result still holds for any admissible Banach $C(\mathcal{X})$ -valued function algebra. But, the following shows that it does hold for admissible $C(\mathcal{X})$ -valued uniform algebras.

Example 4.3. Assume that $A = C(\mathcal{X})$, for some compact Hausdorff space \mathcal{X} . Then, for every admissible A -valued uniform algebra \mathcal{A} on X , the mapping \mathcal{J} in (3.3) is surjective, and thus $\mathfrak{M}(\mathcal{A}) = \mathfrak{M}(\mathfrak{A}) \times \mathfrak{M}(C(\mathcal{X}))$.

To see this, first we show that \mathcal{A} is isometrically isomorphic to $C(\mathcal{X}, \mathfrak{A})$. Take a function $f \in \mathcal{A}$. Then $f(x)$, for every $x \in X$, is a function in $C(\mathcal{X})$. Define $\tilde{f} : \mathcal{X} \rightarrow \mathfrak{A}$ by $\tilde{f}(\xi)(x) = f(x)(\xi)$. In fact, $\tilde{f}(\xi) = \phi_\xi \circ f$ where ϕ_ξ is the evaluation character of $A = C(\mathcal{X})$ at ξ , and, since \mathcal{A} is admissible, $\tilde{f}(\xi)$ belongs to \mathfrak{A} . Now, define $T : \mathcal{A} \rightarrow C(\mathcal{X}, \mathfrak{A})$ by $Tf = \tilde{f}$. It is easily verified that T is an algebra homomorphism, and

$$\|f\|_X = \sup_{x \in X} \|f(x)\| = \sup_{x \in X} \sup_{\xi \in \mathcal{X}} |f(x)(\xi)| = \sup_{\xi \in \mathcal{X}} \|\tilde{f}(\xi)\| = \|\tilde{f}\|_{\mathcal{X}}.$$

Since the range of T contains all elements of the form $g_1 h_1 + \dots + g_n h_n$, where $n \in \mathbb{N}$, $g_i \in C(\mathcal{X})$ and $h_i \in \mathfrak{A}$, and these functions are dense in $C(\mathcal{X}, \mathfrak{A})$, we have T surjective. It follows that T is an isometric isomorphism. By [9, Theorem], $\mathfrak{M}(C(\mathcal{X}, \mathfrak{A}))$ is identical to $\mathfrak{M}(\mathfrak{A}) \times \mathcal{X}$, which means that $\mathfrak{M}(\mathcal{A})$ is identical to $\mathfrak{M}(\mathfrak{A}) \times \mathfrak{M}(A)$.

Example 4.4 (Tensor Products). Let \mathfrak{A} be a Banach function algebra on X and consider the algebraic tensor product $\mathfrak{A} \otimes A$. There exists, by [5, Theorem 42.6], a linear operator $T : \mathfrak{A} \otimes A \rightarrow \mathfrak{A}A$ such that

$$(4.2) \quad T\left(\sum_{i=1}^n f_i \otimes a_i\right) = \sum_{i=1}^n f_i a_i.$$

The operator T is an algebra isomorphism so that $\mathfrak{A} \otimes A$ can be seen as an admissible A -valued function algebra on X . We identify every element

$f \in \mathfrak{A} \otimes A$ with its image Tf as an A -valued function on X . Let $\|\cdot\|_\gamma$ be an algebra cross-norm on $\mathfrak{A} \otimes A$ so that the completion $\mathfrak{A} \widehat{\otimes}_\gamma A$ is a Banach algebra. The mapping T extends to an isometric isomorphism of $\mathfrak{A} \widehat{\otimes}_\gamma A$ onto a Banach A -valued function algebra on X . For example, if $\|\cdot\|_\epsilon$ is the injective tensor norm, then $\mathfrak{A} \widehat{\otimes}_\epsilon A$ is isometrically isomorphic to the uniform closure $\overline{\mathfrak{A}A}$ of $\mathfrak{A}A$ and $\|f\|_\epsilon = \|f\|_X$, for all $f \in \mathfrak{A} \otimes A$.

It is proved in [4] that

- (1) $\mathfrak{A} \widehat{\otimes}_\gamma A$ is an admissible Banach A -valued function algebra on X .
- (2) If $f \in \mathfrak{A} \widehat{\otimes}_\gamma A$ and $\phi \in A^*$ then $\phi \circ f \in \mathfrak{A}$ and $\|\phi \circ f\| \leq \|\phi\| \|f\|_\gamma$.

We now show that every $\tau \in \mathfrak{M}(\mathfrak{A} \widehat{\otimes}_\gamma A)$ is of the form $\tau = \psi \diamond \phi$, with $\phi = \tau|_A$ and $\psi = \tau|_{\mathfrak{A}}$. Since $\mathfrak{A} \otimes A$ is dense in $\mathfrak{A} \widehat{\otimes}_\gamma A$, it is enough to show that $\tau = \psi \diamond \phi$ on $\mathfrak{A} \otimes A$. First, note that every $f \in \mathfrak{A} \otimes A$ can be seen, through the isomorphism (4.2), as $f = f_1 a_1 + \cdots + f_n a_n$. Hence, $\phi \circ f = \phi(a_1) f_1 + \cdots + \phi(a_n) f_n$ and

$$\begin{aligned} \tau(f) &= \tau(f_1 a_1 + \cdots + f_n a_n) \\ &= \tau(f_1) \tau(a_1) + \cdots + \tau(f_n) \tau(a_n) \\ &= \psi(f_1) \phi(a_1) + \cdots + \psi(f_n) \phi(a_n) \\ &= \psi(\phi(a_1) f_1 + \cdots + \phi(a_n) f_n) \\ &= \psi(\phi \circ f). \end{aligned}$$

This proves that $\tau = \psi \diamond \phi$ on $\mathfrak{A} \otimes A$ and thus $\tau = \psi \diamond \phi$ on $\mathfrak{A} \widehat{\otimes}_\gamma A$. We conclude that $\mathfrak{M}(\mathfrak{A} \widehat{\otimes}_\gamma A) = \mathfrak{M}(\mathfrak{A}) \times \mathfrak{M}(A)$. This result, however, can be derived from the following more general result due to Tomiyama [17].

Theorem 4.5 ([17]). *Suppose that A and B are commutative Banach algebras. If $A \widehat{\otimes}_\gamma B$ is a Banach algebra for a cross-norm γ , then $\mathfrak{M}(A \widehat{\otimes}_\gamma B)$ is homeomorphic to $\mathfrak{M}(A) \times \mathfrak{M}(B)$.*

Acknowledgment. The author expresses his sincere gratitude to the referee for their careful reading and suggestions that improved the presentation of this paper.

REFERENCES

- [1] M. Abel, M. Abtahi, *Description of closed maximal ideals in topological algebras of continuous vector-valued functions*, *Mediterr. J. Math.* **11** 4 (2014), 1185–1193.
- [2] M. Abtahi, *Vector-valued characters on vector-valued function algebras*, *Banach J. Math. Anal.* (to appear) [arXiv:1509.09215](https://arxiv.org/abs/1509.09215).
- [3] M. Abtahi, *Normed algebras of infinitely differentiable Lipschitz functions*, *Quaest. Math.* **35** 2 (2012), 195–202.
- [4] M. Abtahi and S. Farhangi, *Vector-valued spectra of Banach algebra valued continuous functions*, (preprint) [arXiv:1510.06641](https://arxiv.org/abs/1510.06641)
- [5] F. F. Bonsall, J. Duncan, *Complete Normed Algebras*, Springer-Verlag, Berlin, Heidelberg, New York, 1973.
- [6] H.G. Dales, *Banach Algebras and Automatic Continuity*. London Mathematical Society Monographs, New Series 24, Oxford Science Publications, Clarendon Press, Oxford University Press, New York, 2000.

- [7] K. Esmaeili and H. Mahyar, *The character spaces and Šilov boundaries of vector-valued lipschitz function algebras*, Indian J. Pure Appl. Math., **45** 6 (2014), 977–988.
- [8] T.W. Gamelin, *Uniform Algebras*, Prentice-Hall, Inc., Englewood Cliffs, NJ., 1969.
- [9] A. Hausner, *Ideals in a certain Banach algebra*, Proc. Amer. Math. Soc. **8** (1957), 246–249.
- [10] T.G. Honary, *Relations between Banach function algebras and their uniform closures*, Proc. Amer. Math. Soc., **109** 2 (1990), 337–342.
- [11] T.G. Honary, A. Nikou and A.H. Sanatpour, *On the character space of vector-valued Lipschitz algebras*, Bull. Iranian Math. Soc. **40** 6 (2014), 1453–1468.
- [12] J. Horváth, *Topological Vector Spaces and Distributions*, Vol. I, Addison-Wesley, Reading, Mass., 1966.
- [13] E. Kaniuth, *A Course in Commutative Banach Algebras*, Graduate Texts in Mathematics, 246 Springer, 2009.
- [14] G.M. Leibowitz, *Lectures on Complex Function Algebras*, Scott, Foresman and Company, 1970.
- [15] A. Nikou and A.G. O’Farrell, *Banach algebras of vector-valued functions*, Glasgow Math. J. **56** 2 (2014), 419–426.
- [16] E.L. Stout, *The Theory of Uniform Algebras*, Bogden & Quigley, Inc., Tarrytown-on-Hudson, New York, 1971.
- [17] J. Tomiyama, *Tensor products of commutative Banach algebras*, Tôhoku Math. J. **12** (1960), 147–154.
- [18] B. Yood, *Banach algebras of continuous functions*, Amer. J. Math., **73** (1951), 30–42.
- [19] W. Żelazko, *Banach Algebras*, Translated from the Polish by Marcin E. Kuczma. Elsevier Publishing Co., Amsterdam-London-New York; PWN–Polish Scientific Publishers, Warsaw, 1973.

MORTAZA ABTAHI

SCHOOL OF MATHEMATICS AND COMPUTER SCIENCES,
DAMGHAN UNIVERSITY, DAMGHAN, P.O.BOX 36715-364, IRAN
E-mail address: abtahi@du.ac.ir; mortaza.abtahi@gmail.com